

At the Cutting Edge

Environmental chemicals as thyroid hormone analogues: New studies indicate that thyroid hormone receptors are targets of industrial chemicals?

R. Thomas Zoeller*

*Biology Department and Program in Molecular and Cellular Biology, Morrill Science Center,
University of Massachusetts-Amherst, Amherst, MA 01003, USA*

Received 22 July 2005; received in revised form 25 July 2005; accepted 27 July 2005

Abstract

Thyroid hormone (TH) is essential for normal brain development, but the specific actions of TH differ across developmental time and brain region. These actions of TH are mediated largely by a combination of thyroid hormone receptor (TR) isoforms that exhibit specific temporal and spatial patterns of expression during animal and human brain development. In addition, TR action is influenced by different co-factors, proteins that directly link the TR protein to functional changes in gene expression. Several recent studies now show that TRs may be unintended targets of chemicals manufactured for industrial purposes, and to which humans and wildlife are routinely exposed. Polychlorinated biphenyls (PCBs), polybrominated diphenyl ethers (PBDEs), and bisphenol-A (BPA), and specific halogenated derivatives and metabolites of these compounds, have been shown to bind to TRs and perhaps have selective effects on TR functions. A number of common chemicals including polybrominated biphenyls (PBBs) and phthalates may also exert such effects. Considering the importance of TH in brain development, it will be important to pursue the possibilities that these chemicals – or interactions among chemical classes – are affecting children’s health by influencing TH signaling in the developing brain.

© 2005 Elsevier Ireland Ltd. All rights reserved.

Keywords: Brain development; Endocrine disruption; Bisphenol A; Poly

Contents

Conclusions	12
References	13

It has long been recognized that thyroid hormone (TH) is essential for normal brain development in both humans and in animals (Dussault and Ruel, 1987), but the mechanisms by which TH exerts its actions are only partially understood (Bernal et al., 2003). Likewise, it has long been recognized that there are environmental influences on thyroid function (Gaitan, 1989), but our ability to identify environmental factors that affect thyroid hormone action during brain development may be limited by the lack of information about thyroid hormone action in the developing brain (Zoeller,

2003). Moreover, because all known “thyroid toxicants” have been identified solely by their ability to reduce circulating TH levels (Brucker-Davis, 1998), the default approach to identify such chemicals is by their effects on hormone levels and on thyroid histology (e.g., size of the colloid, qualitative appearance of hypertrophic or hyperplastic effects) (DeVito et al., 1999). However, chemicals that act directly on thyroid hormone receptors (TRs) may produce variable and perhaps unpredicted effects on hormone levels as well as to produce effects on brain development that incompletely mimic TH insufficiency (or action).

Despite early speculations that environmental chemicals may act as imperfect TH analogues (McKinney and Waller, 1994; McKinney and Waller, 1998), few studies had tested

* Tel.: +1 413 545 2088; fax: +1 413 545 3243.
E-mail address: tzoeller@bio.umass.edu.

this hypothesis until recently. Now, several recent reports show that a broad range of chemicals to which humans are routinely, and inadvertently, exposed can bind to TRs and may produce complex effects on thyroid hormone signaling. Perhaps the best example is that of polychlorinated biphenyls (PCBs), industrial chemicals consisting of paired phenyl rings with various degrees of chlorination (Chana and Concejero, 2002). Although the production of PCBs was banned in the mid 1970's, these contaminants are routinely detected in the environment (Breivik et al., 2002) and in human tissues (Fisher, 1999) at high concentrations. PCB body burden is associated with lower full-scale IQ, reduced visual recognition memory, attention deficits, and motor deficits (Huisman et al., 1995; Jackson et al., 1997; Osius et al., 1999; Korrick and Altshul, 1998; Ayotte et al., 2003; Walkowiak et al., 2001).

PCBs reduce circulating levels of T_4 in animals (Zoeller et al., 2000; Goldey et al., 1995; Bastomsky, 1977a,b; Bastomsky et al., 1976), and some authors propose that PCBs exert neurotoxic effects on the developing brain by causing a state of relative hypothyroidism (Crofton, 2004; Crofton et al., 2000; Brouwer et al., 1999). This concept is supported by the observations that the ototoxic effect of PCB exposure can be partially ameliorated by T_4 replacement (Goldey and Crofton, 1998), and that the cerebellum, a tissue highly sensitive to thyroid hormone insufficiency (Koibuchi and Chin, 2000; Li et al., 2004; Yousefi et al., 2005), is targeted by PCB exposure. PCB exposure alters motor behavior associated with cerebellar function (Roegge et al., 2004; Nguon et al., 2005), as well as cerebellar anatomy (Nguon et al., 2005). Interestingly, PCB exposure is associated with an increase in expression of glial fibrillary acidic protein (GFAP) (Nguon et al., 2005), which is also increased by thyroid hormone insufficiency (Granholt, 1985). Finally, in young children, the association between PCB body burden and behavioral measures of response-inhibition is stronger in those children that have a smaller corpus callosum (Stewart et al., 2003), an area of the brain affected by thyroid hormone (Traggiai and Stanhope, 2004; Schoonover et al., 2004; Berbel et al., 1994). Thus, it is possible that PCBs exert at least some neurotoxic effects on the developing cerebellum by causing a state of relative hypothyroidism.

However, PCB exposure does not produce consistent effects on animals that are indicative of thyroid hormone insufficiency, such as body weight gain during development (Zoeller et al., 2000; Gauger et al., 2004; Bansal et al., 2005) or the timing of eye opening (Goldey et al., 1995). In addition, despite the reduction in serum T_4 , PCB exposure increases the expression of several thyroid hormone-responsive genes in the fetal (Gauger et al., 2004; Bansal et al., 2005) and neonatal (Zoeller et al., 2000) brain. These observations are consistent with the hypothesis that at least some individual PCB congeners, or their metabolites, can act as TR agonists in vivo. Recently, Kitamura et al. (2005) reported on their observations that nine separate hydroxylated PCB congeners can bind to the rat TR with an

IC_{50} as low as 5 μ M. In addition, using a human neuroprogenitor cell line, Fritsche et al. (2005) found that a specific PCB congener could mimic the ability of T_3 in increasing oligodendrocyte differentiation, and that this effect was blocked by the selective TR antagonist NH3. Finally, Arulmozhiraja and Morita (2004) have identified several PCB congeners that exhibit weak thyroid hormone activity in a yeast-two hybrid assay optimized to identify such activity.

However, not all recent reports indicate that PCBs act as agonists on the TR. Kimura-Kuroda et al. (2005) have found that two separate hydroxylated PCBs interfere with T_3 -dependent neurite outgrowth in mouse cerebellar granule cell primary cultures. In addition, Bogazzi et al. (2003) found that a commercial mixture of PCBs (Aroclor 1254) exhibited specific binding to the rat TR β at approximately 10 μ M. This concentration inhibited TR action on the malic enzyme promoter in a CAT assay and this effect required an intact TRE. However, the PCB mixture did not alter the ability of TR to bind to the ME TRE in a gel shift assay. In contrast, Iwasaki et al. (2002) found that a specific hydroxylated PCB congener inhibits TR-mediated transcriptional activation in a luciferase assay at concentrations as low as 10^{-10} M. This effect was observed in several cell lines, but was not observed using a glucocorticoid response element. Miyazaki et al. (2004) followed this report by showing that PCBs can dissociate TR:RXR heterodimers from a TRE.

It is clear that PCBs are neurotoxic in humans and animals, and that they can interact directly with the TR. However, the consequences of PCB exposure on TR action appear to be quite complex. This complexity includes acting as an agonist or antagonist and may include TR isoform selectivity in as much as most studies have been performed using the TR β , leaving the TR α relatively unstudied in this context. In addition, considering that there are 209 different chlorine substitution patterns on the biphenyl backbone and that these can be metabolized (hydroxyl- and methylsulfonyl-metabolites (Kato et al., 1998)), it is possible that different chemical species exerts different effects. Finally, PCBs may exert different actions on TRs depending on associated heterodimer partners, promoter structure, or different co-factors. This complexity will be important to pursue because the effects of PCB exposure in humans is far better studied than for structurally related compounds such as polybrominated biphenyls (PBBs) and polybrominated diphenyl ethers (PBDEs). Thus, mechanistic studies on PCBs can be more easily and effectively coupled to specific human health outcomes.

Bisphenol-A (BPA, 4,4' isopropylidenediphenol) is produced at a rate of over 800 million kg annually in the United States alone (Reporter, 1999), and is used primarily in the manufacture of plastics including polycarbonate plastics, epoxy resins that coat food cans, and in dental sealants (Howe et al., 1998; Lewis et al., 1999). Howe et al. (1998) estimated human consumption of BPA from epoxy-lined food cans alone to be about 6.6 μ g/person-day. BPA has been reported in concentrations of 1–10 ng/ml in serum of pregnant women, in the amniotic fluid of their fetus, and in cord serum taken at

birth (Schonfelder et al., 2002; Ikezuki et al., 2002). Moreover, BPA concentrations of up to 100 ng/g were reported in placenta (Schonfelder et al., 2002). Bisphenol-A is also halogenated (brominated or chlorinated) to produce flame-retardants. Tetrabromobisphenol-A (TBBPA) is the most commonly used with over 60,000 tons produced annually (WHO, 1995; WHO, 1997). Thomsen et al. (2002) recently reported that brominated flame retardants, including TBBPA, have increased in human serum from 1977 to 1999 with concentrations in adults ranging from 0.4 to 3.3 ng/g serum lipids. However, infants (0–4 years) exhibited serum concentrations that ranged from 1.6 to 3.5 times higher (Thomsen et al., 2002).

Considering this pattern of human exposure, it is potentially important that BPA has been shown to bind to the TR (Moriyama et al., 2002a). Best characterized as a weak estrogen (Staples et al., 1998), binding to the estrogen receptor (ER) with a K_i of approximately 10^{-5} M (Krishnan et al., 1993; Gaido et al., 1997), BPA binds to and antagonizes T_3 activation of the TR (Kitamura et al., 2002; Moriyama et al., 2002b) with a K_i of approximately 10^{-4} M, but as little as 10^{-6} M BPA significantly inhibits TR-mediated gene activation (Moriyama et al., 2002b). Moreover, Moriyama et al. (2002a) found that BPA reduced T_3 -mediated gene expression in culture by enhancing the interaction with the co-repressor N-CoR. Interestingly; we have found that developmental exposure to BPA in rats produces an endocrine profile similar to that observed in thyroid resistance syndrome (Cheng, 2005). Specifically, T_4 levels were elevated during development in the pups of BPA-treated animals, but TSH levels were not different from controls (Zoeller et al., 2005). This profile is consistent with BPA inhibition of $TR\beta$ -mediated negative feedback. However, the thyroid hormone-response gene RC3 was elevated in the dentate gyrus of these BPA-treated animals (Zoeller et al., 2005). Because the $TR\alpha$ isoform is expressed in the dentate gyrus, we concluded that BPA may be a selective $TR\beta$ antagonist in vivo.

If BPA acts as a TR antagonist in vivo, it is predictable that specific developmental events and behaviors would be affected by developmental exposure to BPA. In this regard, Seiwa et al. (2004) have shown that BPA blocks T_3 -induced oligodendrocyte development from precursor cells (OPCs). In addition, there may be an association between the thyroid resistance syndrome and attention deficit-hyperactivity disorder (ADHD) in humans (Siesser et al., 2005; Vermiglio et al., 2004; Hauser et al., 1998) and in rats (Siesser et al., 2005) therefore, it is potentially important that BPA-exposed rats exhibit ADHD-like symptoms (Ishido et al., 2004).

Despite the antagonistic effects of BPA on the $TR\beta$, halogenated BPAs appear to act as TR agonists (Kitamura et al., 2002). Both tetrabromo and tetrachlorobisphenol A (TBBPA and TCBPA, respectively) can bind to the thyroid hormone receptor and induce GH3 cell proliferation and growth hormone production (Kitamura et al., 2002). Thus, these compounds may exert agonistic effects on the TR and this could be




Fig. 1. The current theory of thyroid hormone action on the TR allows us to predict sites at which environmental chemicals may interfere with thyroid hormone action. Chemicals may bind to the TR and directly activate or inhibit the action of endogenous T_3 (1). This action may occur by influencing the interaction of TRs with various co-factors such as N-CoR or SRC-1 (2). In addition, environmental chemicals may cause the TR to exhibit a different affinity for the TRE (3). These effects may well be dependent upon the specific TR isoform ($TR\alpha$ or $TR\beta$ isoforms), the specific TRE or contextual sequences of specific TREs, and/or the specific co-factors available in the cell. It seems predictable that these chemicals will not produce patterns of effects or disease that simply mimic thyroid hormone insufficiency or excess and thus may easily be misinterpreted both in experimental animals and in humans.

important during early brain development. For example, thyroid hormone of maternal origin can regulate gene expression in the fetal brain (Dowling et al., 2000,2001; Dowling and Zoeller, 2000); one of these genes codes for Hes1 (Bansal et al., 2005). Considering the role of HES proteins in fate specification in the early cortex (Wu et al., 2003; Schuurmans and Guillemot, 2002; Gaiano and Fishell, 2002), the observation that industrial chemicals can activate the TR and increase HES expression (Bansal et al., 2005) may indicate that these chemicals can exert subtle effects on early differentiative events.

Conclusions

The human population is exposed to a large number of specific polyhalogenated aromatic hydrocarbons, and biomonitoring studies now detect these chemicals in adults, children, pregnant women and in the fetal compartment (Takser et al., 2005). Increasing numbers of reports are revealing that a broad array of compounds can bind to the TR and affect TH-regulated gene expression, both in vivo and in vitro. However, considering the tremendously pleiotropic effects of thyroid hormone, it is predictable that these synthetic compounds may have very complex effects on the TR (Fig. 1). In addition, these studies suggest that chemicals may interact with other important TH binding proteins. For example, deiodinase enzymes appear to control the sensitivity of different brain regions to TH exposure during development (Kester et al., 2004); thus, if exogenous chemicals alter the activity of these enzymes, it may influence the sequence of TH-sensitive developmental events. Likewise, specific transporters appear to control the availability of T_3 to cells in the brain (Friesema et al., 2003; Heuer et al., 2005); thus, if environmental chemicals interfere with tissue uptake of TH, adverse human health effects could result. Our ability to identify chemical effects

on TR function in vitro far exceeds our ability to identify chemical effects on TR function in vivo, in part because the mechanisms of TH action in the developing brain is less well understood. However, it will be important to define the role of TH in brain development and to identify the mechanisms by which TH exerts these actions if we are to understand the potential human health effects of persistent exposure to these bioaccumulative compounds.

References

- Arulmozhiraja, S., Morita, M., 2004. Structure-activity relationships for the toxicity of polychlorinated dibenzofurans: approach through density functional theory-based descriptors. *Chem. Res. Toxicol.* 17, 348–356.
- Ayotte, P., Muckle, G., Jacobson, J.L., Jacobson, S.W., Dewailly, E., 2003. Assessment of pre- and postnatal exposure to polychlorinated biphenyls: lessons from the inuit cohort study. *Environ. Health Perspect.* 111, 1253–1258.
- Bansal, R., You, S.H., Herzig, C.T., Zoeller, R.T., 2005. Maternal thyroid hormone increases HES expression in the fetal rat brain: an effect mimicked by exposure to a mixture of polychlorinated biphenyls (PCBs). *Brain Res. Dev. Brain Res.* 156, 13–22.
- Bastomsky, C.H., Murthy, P.V.N., Banovac, K., 1976. Alterations in thyroxine metabolism produced by cutaneous application of microscope immersion oil: effects due to polychlorinated biphenyls. *Endocrinology* 98, 1309–1314.
- Bastomsky, C.H., 1977a. Goitres in rats fed polychlorinated biphenyls. *Can. J. Physiol. Pharmacol.* 55, 288–292.
- Bastomsky, C.H., 1977b. Enhanced thyroxine metabolism and high uptake goiters in rats after a single dose of 2,3,7,8-tetrachlorodibenzo-*p*-dioxin. *Endocrinology* 101, 292–296.
- Berbel, P., Guadano-Ferraz, A., Angulo, A., Raman Cerezo, J., 1994. Role of thyroid hormones in the maturation of interhemispheric connections in rats. *Behav. Brain Res.* 64, 9–14.
- Bernal, J., Guadano-Ferraz, A., Morte, B., 2003. Perspectives in the study of thyroid hormone action on brain development and function. *Thyroid* 13, 1005–1012.
- Bogazzi, F., Raggi, F., Ultimieri, F., Russo, D., Campomori, A., McKinney, J.D., Pinchera, A., Bartalena, L., Martino, E., 2003. Effects of a mixture of polychlorinated biphenyls (Aroclor 1254) on the transcriptional activity of thyroid hormone receptor. *J. Endocrinol. Invest.* 26, 972–978.
- Brevik, K., Sweetman, A., Pacyna, J.M., Jones, K.C., 2002. Towards a global historical emission inventory for selected PCB congeners—a mass balance approach. 1. Global production and consumption. *Sci. Total Environ.* 290, 181–198.
- Brouwer, A., Longnecker, M.P., Birnbaum, L.S., Cogliano, J., Kostyniak, P., Moore, J., Schantz, S., Winneke, G., 1999. Characterization of potential endocrine-related health effects at low-dose levels of exposure to PCBs. *Environ. Health Perspect.* 107 (Suppl. 4), 639–649.
- Brucker-Davis, F., 1998. Effects of environmental synthetic chemicals on thyroid function. *Thyroid* 8, 827–856.
- Chana, A., Concejero, M.A., de Frutos, M.A., Gonzalez, M.J., Herradon, B., 2002. Computational studies on biphenyl derivatives. Analysis of the conformational mobility, molecular electrostatic potential, and dipole moment of chlorinated biphenyl: searching for the rationalization of the selective toxicity of polychlorinated biphenyls (PCBs). *Chem. Res. Toxicol.* 15, 1514–1526.
- Cheng, S.Y., 2005. Thyroid hormone receptor mutations and disease: beyond thyroid hormone resistance. *Trends Endocrinol. Metab.* 16, 176–182.
- Crofton, K.M., Kodavanti, P.R., Derr-Yellin, E.C., Casey, A.C., Kehn, L.S., 2000. PCBs, thyroid hormones, and ototoxicity in rats: cross-fostering experiments demonstrate the impact of postnatal lactation exposure. *Toxicol. Sci.* 57, 131–140.
- Crofton, K.M., 2004. Developmental disruption of thyroid hormone: correlations with hearing dysfunction in rats. *Risk Anal.* 24, 1665–1671.
- DeVito, M., Biegel, L., Brouwer, A., Brown, S., Brucker-Davis, F., Cheek, A.O., Christensen, R., Colborn, T., Cooke, P., Crissman, J., Crofton, K., Doerge, D., Gray, E., Hauser, P., Hurley, P., Kohn, M., Lazar, J., McMaster, S., McClain, M., McConnell, E., Meier, C., Miller, R., Tietge, J., Tyl, R., 1999. Screening methods for thyroid hormone disruptors. *Environ. Health Perspect.* 107, 407–415.
- Dowling, A.L.S., Zoeller, R.T., 2000. Thyroid hormone of maternal origin regulates the expression of RC3/Neurogranin mRNA in the fetal rat brain. *Brain Res.* 82, 126–132.
- Dowling, A.L.S., Martz, G.U., Leonard, J.L., Zoeller, R.T., 2000. Acute changes in maternal thyroid hormone induce rapid and transient changes in specific gene expression in fetal rat brain. *J. Neurosci.* 20, 2255–2265.
- Dowling, A.L.S., Iannacone, E.A., Zoeller, R.T., 2001. Maternal hypothyroidism selectively affects the expression of neuroendocrine-specific protein-A messenger ribonucleic acid in the proliferative zone of the fetal rat brain cortex. *Endocrinology* 142, 390–399.
- Dussault, J.H., Ruel, J., 1987. Thyroid hormones and brain development. *Annu. Rev. Physiol.* 49, 321–334.
- Fisher, B.E., 1999. Most unwanted. *Environ. Health Perspect.* 107, A18–A23.
- Friesema, E.C., Ganguly, S., Abdalla, A., Manning Fox, J.E., Halestrap, A.P., Visser, T.J., 2003. Identification of monocarboxylate transporter 8 as a specific thyroid hormone transporter. *J. Biol. Chem.* 278, 40128–40135.
- Fritsche, E., Cline, J.E., Nguyen, N.H., Scanlan, T.S., Abel, J., 2005. Polychlorinated biphenyls disturb differentiation of normal human neural progenitor cells: clue for involvement of thyroid hormone receptors. *Environ. Health Perspect.* 113, 871–876.
- Gaiano, N., Fishell, G., 2002. The role of notch in promoting glial and neural stem cell fates. *Annu. Rev. Neurosci.* 25, 471–490.
- Gaido, K.W., Leonard, L.S., Lovell, S., Gould, J.C., Babai, D., Portier, C.J., McDonnell, D.P., 1997. Evaluation of chemicals with endocrine modulating activity in yeast-based steroid hormone receptor gene transcription assay. *Toxicol. Appl. Pharmacol.* 143, 205–212.
- Gaitan, E. (Ed.), 1989. *Environmental Goitrogenesis*. CRC Press, Inc., Boca Raton, p. 250.
- Gauger, K.J., Kato, Y., Haraguchi, K., Lehmler, H.J., Robertson, L.W., Bansal, R., Zoeller, R.T., 2004. Polychlorinated biphenyls (PCBs) exert thyroid hormone-like effects in the fetal rat brain but do not bind to thyroid hormone receptors. *Environ. Health Perspect.* 112, 516–523.
- Goldey, E.S., Crofton, K.M., 1998. Thyroxine replacement attenuates hypothyroxinemia, hearing loss, and motor deficits following developmental exposure to Aroclor 1254 in rats. *Toxicol. Sci.* 45, 94–105.
- Goldey, E.S., Kehn, L.S., Lau, C., Rehnberg, G.L., Crofton, K.M., 1995. Developmental exposure to polychlorinated biphenyls (Aroclor 1254) reduces circulating thyroid hormone concentrations and causes hearing deficits in rats. *Toxicol. Appl. Pharmacol.* 135, 77–88.
- Granhölm, A.C., 1985. Effects of thyroid hormone deficiency on glial constituents in developing cerebellum of the rat. *Exp. Brain Res.* 59, 451–456.
- Hauser, P., McMillin, J.M., Bhatara, V.S., 1998. Resistance to thyroid hormone: implications for neurodevelopmental research on the effects of thyroid hormone disruptors. *Toxicol. Ind. Health* 14, 85–101.
- Heuer, H., Maier, M.K., Iden, S., Mittag, J., Friesema, E.C., Visser, T., Bauer, K., 2005. The monocarboxylate transporter 8 linked to human psychomotor retardation is highly expressed in thyroid hormone sensitive neuron populations. *Endocrinology* 146, 1701–1706.
- Howe, S.R., Borodinsky, L., Lyon, R.S., 1998. Potential exposure to bisphenol A from food-contact use of epoxy coated cans. *J. Coat. Technol.* 70, 69–74.

- Huisman, M., Koopman-Esseboom, C., Lanting, C.I., van der Paauw, C.G., Tuinstra, L.G., Fidler, V., Weisglas-Kuperus, N., Sauer, P.J., Boersma, E.R., Touwen, B.C., 1995. Neurological condition in 18-month-old children perinatally exposed to polychlorinated biphenyls and dioxins. *Early Hum. Dev.* 43, 165–176.
- Ikezuki, Y., Tsutsumi, O., Takai, Y., Kamei, Y., Taketani, Y., 2002. Determination of bisphenol A concentrations in human biological fluids reveals significant early prenatal exposure. *Hum. Reprod.* 17, 2839–2841.
- Ishido, M., Masuo, Y., Kunimoto, M., Oka, S., Morita, M., 2004. Bisphenol A causes hyperactivity in the rat concomitantly with impairment of tyrosine hydroxylase immunoreactivity. *J. Neurosci. Res.* 76, 423–433.
- Iwasaki, T., Miyazaki, W., Takeshita, A., Kuroda, Y., Koibuchi, N., 2002. Polychlorinated biphenyls suppress thyroid hormone-induced transactivation. *Biochem. Biophys. Res. Commun.* 299, 384–388.
- Jackson, T.A., Richer, J.K., Bain, D.L., Takimoto, G.S., Tung, L., Horwitz, K.B., 1997. The partial agonist activity of antagonist-occupied steroid receptors is controlled by a novel hinge domain-binding coactivator L7/SPA and the corepressors N-CoR or SMRT. *Mol. Endocrinol.* 11, 693–705.
- Kato, Y., Haraguchi, K., Shibahara, T., Masuda, Y., Kimura, R., 1998. Reduction of thyroid hormone levels by methylsulfonyl metabolites of polychlorinated biphenyl congeners in rats. *Arch. Toxicol.* 72, 541–544.
- Kester, M.H.A., Martinez de Mena, R., Obregon, M.J., Marinkovic, D., Visser, T.J., Hume, R., Morreale de Escobar, G., 2004. Iodothyronine levels in the human developing brain: major regulatory roles of iodothyronine deiodinases in different areas. *J. Clin. Endocrinol. Metab.* 89, 3117–3128.
- Kimura-Kuroda, J., Nagata, I., Kuroda, Y., 2005. Hydroxylated metabolites of polychlorinated biphenyls inhibit thyroid-hormone-dependent extension of cerebellar Purkinje cell dendrites. *Brain Res. Dev. Brain Res.* 154, 259–263.
- Kitamura, S., Jinno, N., Ohta, S., Kuroki, H., Fujimoto, N., 2002. Thyroid hormonal activity of the flame retardants tetrabromobisphenol A and tetrachlorobisphenol A. *Biochem. Biophys. Res. Commun.* 293, 554–559.
- Kitamura, S., Jinno, N., Suzuki, T., Sugihara, K., Ohta, S., Kuroki, H., Fujimoto, N., 2005. Thyroid hormone-like and estrogenic activity of hydroxylated PCBs in cell culture. *Toxicology* 208, 377–387.
- Koibuchi, N., Chin, W.W., 2000. Thyroid hormone action and brain development. *Trends Endocrinol. Metab.* 11, 123–128.
- Korrick, S.A., Altshul, L., 1998. High breast milk levels of polychlorinated biphenyls (PCBs) among four women living adjacent to a PCB-contaminated waste site. *Environ. Health Perspect.* 106, 513–518.
- Krishnan, A.V., Stathis, P., Permuth, S.F., Tokes, L., Feidman, D., 1993. Bisphenol A: an estrogenic substance is released from polycarbonate flasks during autoclaving. *Endocrinology* 132, 2279–2286.
- Lewis, J.B., Rueggeberg, F.A., Lapp, C.A., Ergle, J.W., Schuster, G.S., 1999. Identification and characterization of estrogen-like components in commercial resin-based dental restorative materials. *Clin. Oral Invest.* 3, 107–113.
- Li, G.H., Post, J., Koibuchi, N., Sajdel-Sulkowska, E.M., 2004. Impact of thyroid hormone deficiency on the developing CNS: cerebellar glial and neuronal protein expression in rat neonates exposed to antithyroid drug propylthiouracil. *Cerebellum* 3, 100–106.
- McKinney, J.D., Waller, C.L., 1994. Polychlorinated biphenyls as hormonally active structural analogues. *Environ. Health Perspect.* 102, 290–297.
- McKinney, J.D., Waller, C.L., 1998. Molecular determinants of hormone mimicry: halogenated aromatic hydrocarbon environmental agents. *J. Toxicol. Environ. Health B Crit. Rev.* 1, 27–58.
- Miyazaki, W., Iwasaki, T., Takeshita, A., Kuroda, Y., Koibuchi, N., 2004. Polychlorinated biphenyls suppress thyroid hormone receptor-mediated transcription through a novel mechanism. *J. Biol. Chem.* 279, 18195–18202.
- Moriyama, K., Tagami, T., Akamizu, T., Usui, T., Saijo, M., Kanamoto, N., Hataya, Y., Shimatsu, A., Kuzuya, H., Nakao, K., 2002a. Thyroid hormone action is disrupted by bisphenol A as an antagonist. *J. Clin. Endocrinol. Metab.* 87, 5185–5190.
- Moriyama, K., Tagami, T., Akamizu, T., Usui, T., Saijo, M., Kanamoto, N., Hataya, Y., Shimatsu, A., Kuzuya, H., Nakao, K., 2002b. Thyroid hormone action is disrupted by bisphenol A as an antagonist. *J. Clin. Endocrinol. Metab.* 87, 5185–5190.
- Nguon, K., Baxter, M.G., Sajdel-Sulkowska, E.M., 2005. Perinatal exposure to polychlorinated biphenyls differentially affects cerebellar development and motor functions in male and female rat neonates. *Cerebellum* 4, 112–122.
- Osius, N., Karmaus, W., Kruse, H., Witten, J., 1999. Exposure to polychlorinated biphenyls and levels of thyroid hormones in children. *Environ. Health Perspect.* 107, 843–849.
- Reporter, C.M., 1999. ChemExpo Chemical Profile: Bisphenol-A, vol. 2001, Chemical Market Reporter. Schnell Publishing Company.
- Roegge, C.S., Wang, V.C., Powers, B.E., Klintoova, A.Y., Villareal, S., Greenough, W.T., Schantz, S.L., 2004. Motor impairment in rats exposed to PCBs and methylmercury during early development. *Toxicol. Sci.* 77, 315–324.
- Schonfelder, G., Wittfoht, W., Hopp, H., Talsness, C.E., Paul, M., Chahoud, I., 2002. Parent bisphenol A accumulation in the human maternal-fetal-placental unit. *Environ. Health Perspect.* 110, A703–A707.
- Schoonover, C.M., Seibel, M.M., Jolson, D.M., Stack, M.J., Rahman, R.J., Jones, S.A., Mariash, C.N., Anderson, G.W., 2004. Thyroid hormone regulates oligodendrocyte accumulation in developing rat brain white matter tracts. *Endocrinology* 145, 5013–5020.
- Schuermans, C., Guillemot, F., 2002. Molecular mechanisms underlying cell fate specification in the developing telencephalon. *Curr. Opin. Neurobiol.* 12, 26–34.
- Seiwa, C., Nakahara, J., Komiyama, T., Katsu, Y., Iguchi, T., Asou, H., 2004. Bisphenol A exerts thyroid-hormone-like effects on mouse oligodendrocyte precursor cells. *Neuroendocrinology* 80, 21–30.
- Siesser, W.B., Cheng, S.Y., McDonald, M.P., 2005. Hyperactivity, impaired learning on a vigilance task, and a differential response to methylphenidate in the TRbetaPV knock-in mouse. *Psychopharmacology (Berl)*, 1–11.
- Staples, C.A., Dorn, P.B., Klecka, G.M., O'Block, S.T., Harris, L.R., 1998. A review of the environmental fate, effects, and exposures of bisphenol A. *Chemosphere* 36, 2149–2173.
- Stewart, P., Fitzgerald, S., Reihman, J., Gump, B., Lonky, E., Darvill, T., Pagano, J., Hauser, P., 2003. Prenatal PCB exposure, the corpus callosum, and response inhibition. *Environ. Health Perspect.* 111, 1670–1677.
- Takser, L., Mergler, D., Baldwin, M., de Grosbois, S., Smargiassi, A., Lafond, J., 2005. Thyroid hormones in pregnancy in relation to environmental exposure to organochlorine compounds and mercury. *Environ. Health Perspect.* 113, 1039–1045.
- Thomsen, C., Lundanes, E., Becher, G., 2002. Brominated flame retardants in archived serum samples from Norway: a study on temporal trends and the role of age. *Environ. Sci. Technol.* 36, 1414–1418.
- Traggiai, C., Stanhope, R., 2004. Body mass index and hypothalamic morphology on MRI in children with congenital midline cerebral abnormalities. *J. Pediatr. Endocrinol. Metab.* 17, 219–221.
- Vermiglio, F., Lo Presti, V.P., Moleti, M., Sidoti, M., Tortorella, G., Scafidi, G., Castagna, M.G., Mattina, F., Violi, M.A., Crisa, A., Artemisia, A., Trimarchi, F., 2004. Attention deficit and hyperactivity disorders in the offspring of mothers exposed to mild-moderate iodine deficiency: a possible novel iodine deficiency disorder in developed countries. *J. Clin. Endocrinol. Metab.* 89, 6054–6060.
- Walkowiak, J., Wiener, J.A., Fastabend, A., Heinzow, B., Kramer, U., Schmidt, E., Steingruber, H.J., Wundram, S., Winneke, G., 2001. Environmental exposure to polychlorinated biphenyls and quality of the

- home environment: effects on psychodevelopment in early childhood. *Lancet* 358, 1602–1607.
- WHO, E.H.C., 1995. Tetrabromobisphenol A and Derivatives. World Health Organization, Geneva, Switzerland.
- WHO, E.H.C., 1997. Flame-retardants: a general introduction. World Health Organization, Geneva, Switzerland.
- Wu, Y., Liu, Y., Levine, E.M., Rao, M.S., 2003. Hes1 but not Hes5 regulates an astrocyte versus oligodendrocyte fate choice in glial restricted precursors. *Dev. Dyn.* 226, 675–689.
- Yousefi, B., Jingu, H., Ohta, M., Umezu, M., Koibuchi, N., 2005. Postnatal changes of steroid receptor coactivator-1 immunoreactivity in rat cerebellar cortex. *Thyroid* 15, 314–319.
- Zoeller, R.T., Dowling, A.L., Vas, A.A., 2000. Developmental exposure to polychlorinated biphenyls exerts thyroid hormone-like effects on the expression of RC3/neurogranin and myelin basic protein messenger ribonucleic acids in the developing rat brain. *Endocrinology* 141, 181–189.
- Zoeller, R.T., Bansal, R., Parris, C., 2005. Bisphenol A, an environmental contaminant that acts as a thyroid hormone receptor antagonist in vitro, increases serum thyroxine, and alters RC3/neurogranin expression in the developing rat brain. *Endocrinology* 146, 607–612.
- Zoeller, R.T., 2003. Thyroid toxicology and brain development: should we think differently? *Environ. Health Perspect.* 111, A628.